

Infra-Red Stable Fixed Points of Yukawa Couplings in Supersymmetric Models

B. C. Allanach¹ and S. F. King²

1. Rutherford Appleton Laboratory, Chilton, Didcot, OX11 0QX, U.K.
2. Department of Physics and Astronomy, University of Southampton, Southampton, SO17 1BJ, U.K.

Abstract

We provide the solutions of the fixed point conditions of the Yukawa sector for a large class of $N = 1$ supersymmetric theories including the minimal and next-to-minimal supersymmetric standard models and their grand unified and other extensions. We also introduce a test which can discriminate between infra-red stable, infra-red unstable and saddle point solutions, and illustrate our methods with the example of the next-to-minimal supersymmetric standard model. We show that in this case, the fixed point prediction of the top quark mass is equivalent to that of the minimal supersymmetric standard model, supporting previous numerical analyses.

The fermion mass problem in the standard model arises from the presence of many unknown dimensionless Yukawa couplings. A possible solution to this long-standing problem is that the standard model could be embedded in some more predictive high energy theory with fewer Yukawa couplings. Alternatively one may attempt to relate the Yukawa couplings to the gauge coupling, as in the Pendleton-Ross infra-red stable fixed point (IRSFP) for the top quark Yukawa coupling [1], or the quasi-fixed point of Hill [2]¹. To obtain IRSFPs it is necessary that the unknown dimensionless Yukawa couplings are of the same order as the gauge coupling at high energy. Thus at first sight this approach would seem to be inapplicable to the small Yukawa couplings of the standard model. However string theory yields large Yukawa couplings of order the gauge coupling, and this could suggest that the small dimensionless Yukawa couplings be reinterpreted as large Yukawa couplings multiplied by a ratio of mass scales. In such a scenario fixed points may once again be applicable, as pointed out recently by Ross [4]. In this case the high energy scale would be the string scale and the infra-red region would be near the GUT scale. Thus IRSFPs may provide an unexpected resolution of the fermion mass problem. IRSFPs in the MSSM can also be used to explain the apparent success of the $\lambda_\tau = \lambda_b$ GUT scale relation of some popular SUSY GUTs [5]. Further fixed points have also been identified in the soft masses of supersymmetric theories [6, 7, 8]. It is clear that the idea of IRSFPs has general applicability, and here we shall follow a model independent approach.

In this letter we shall present a general approach which may be used to identify the fixed points that exist in the dimensionless Yukawa couplings of a large class of $N = 1$ supersymmetric theories, and to determine the stability properties of the fixed points in the infra-red. Our assumptions are listed below:

1. We work at one loop order only.
2. We assume that the wavefunction anomalous dimensions are diagonal $\gamma_i^j \propto \delta_i^j \gamma_i$. This may be due to some symmetry that forbids wavefunction mixing between different irreducible representations. For example in the MSSM R-parity forbids mixing between the lepton doublet L and the Higgs doublet H_1 .
3. We assume a single gauge coupling. This may either be due to several approximately equal gauge couplings, which could be the case near the string scale, or due to there being a single dominant gauge coupling as in the case of QCD at low energies where the electroweak couplings may be neglected.

Our general approach may be compared to that of Lanzagorta and Ross [3] for a single Yukawa coupling, however unlike these authors we shall present a well defined procedure for obtaining the fixed points for theories involving any number of large Yukawa couplings in the theory, and having identified the fixed points we shall give a general discussion of their infra-red stability.

¹The relation between the fixed point and quasi-fixed point has been fully elucidated by Lanzagorta and Ross [3].

We shall be interested in the one-loop renormalisation group equations (RGEs) for the dimensionless trilinear Yukawa couplings of an N=1 SUSY theory, which are by now well known and have been widely reported (see for example Martin and Vaughn [9]). The trilinear part of the superpotential may be written

$$W = \frac{1}{6} h^{ijk} \phi_i \phi_j \phi_k, \quad (1)$$

where h^{ijk} are the dimensionless Yukawa couplings, ϕ_i represent the superfields and the sums over i, j, k run over all superfields in the spectrum of the effective theory. The relevant RGEs read

$$16\pi^2 \frac{dh^{ijk}}{d \ln \mu} = h^{ijp} \gamma_p^k + h^{ipk} \gamma_p^j + h^{pjk} \gamma_p^i, \quad (2)$$

$$16\pi^2 \frac{dg_a}{d \ln \mu} = b_a g_a^3 \quad (3)$$

where μ is the $\overline{MS}$ renormalisation scale and the anomalous dimension is defined as

$$\gamma_i^j \equiv \frac{1}{2} \sum_{p,q} h_{ipq} h^{jpq} - 2\delta_i^j g_a^2 C_i^a \quad (4)$$

to one loop order. b_a is the one-loop beta function of the gauge group G_a and the sum over a here represents a sum over all dominant gauge couplings. $C^a(i)$ is the quadratic Casimir of the representation of ϕ_i under the group with gauge coupling g_a :

$$(T_R^a T_R^a)_i^j \equiv C_R^a \delta_i^j. \quad (5)$$

Here, T_R^a is a matrix in the R representation for the group labeled by a . Pictorially, the RG evolution of the trilinear coupling can be described as an insertion of the anomalous dimension correction on each external leg [10].

According to our second assumption, the wavefunction anomalous dimensions are diagonal,

$$\gamma_i^j = \gamma_i \delta_i^j \quad (6)$$

then the above equations become:

$$16\pi^2 \frac{dh^{ijk}}{d \ln \mu} = h^{ijk} (\gamma_i + \gamma_j + \gamma_k), \quad (7)$$

where the anomalous dimension is defined as

$$\gamma_i \equiv \frac{1}{2} \sum_{p,q} h_{ipq} h^{ipq} - 2g_a^2 C_i^a \quad (8)$$

We now introduce a more convenient notation:

$$Y^{ijk} \equiv \frac{(h^{ijk})^2}{16\pi^2}, \quad \tilde{\alpha}_a \equiv \frac{g_a^2}{16\pi^2}, \quad N_i \equiv -\frac{\gamma_i}{16\pi^2}, \quad t \equiv -\ln \mu^2 \quad (9)$$

in terms of which:

$$\frac{dY^{ijk}}{dt} = Y^{ijk}(N_i + N_j + N_k). \quad (10)$$

According to our third assumption there is a single gauge coupling in the theory. We define the ratio of each Yukawa coupling to the single gauge coupling α as:

$$R^{ijk} = \frac{Y^{ijk}}{\tilde{\alpha}} \quad (11)$$

The RGEs now become:

$$\frac{dR^{ijk}}{dt} = \tilde{\alpha} R^{ijk}(N_i + N_j + N_k + \tilde{\alpha}b), \quad (12)$$

The non-trivial (non-zero) IRSFPs we seek correspond to $\frac{dR^{ijk}}{dt} = 0$ for all i, j, k with $R^{ijk} \neq 0$, which implies:

$$N_i + N_j + N_k + \tilde{\alpha}b = 0, \quad (13)$$

for all i, j, k corresponding to $R^{ijk} \neq 0$.

In the above notation the anomalous dimension N_i refers to the superfield ϕ_i while the trilinear coupling between three such superfields is written as a three index tensor h^{ijk} . In a given theory each superfield will in general be in some representation of the gauge group, and i runs over all gauge indices as well as the different representations themselves. Thus most of the entries of the tensor h^{ijk} will be zero due to gauge invariance and other imposed symmetries, and many of the non-zero entries will be equal by gauge invariance. For example in a toy theory with a single top quark Yukawa coupling, assuming that QCD is the only gauge group, the index i would run over the 6 components of the quark doublet $Q = (t, b)$, the 3 components of the top field t^c and the 2 components of the Higgs doublet H , and in general the Yukawa coupling would be h^{ijk} which would be a three index tensor with each index taking 11 possible values. The top quark Yukawa coupling is characterised by a single parameter h , as is clear by dropping all indices and writing the superpotential $W = hQt^cH_2$. In looking for IRSFPs we are interested in the value of h , so it is clear that we must simplify the notation in order to remove all the redundant information contained in the tensor h^{ijk} to arrive at the physical coupling h . In the case of the top quark Yukawa coupling it is clear that we can define $Y_h \equiv h^2/(16\pi^2)$ and write the RGE for the single physical top quark Yukawa coupling as

$$\frac{dY_h}{dt} = Y_h(N_Q + N_{t^c} + N_{H_2}) \quad (14)$$

or defining $R_h \equiv Y_h/\tilde{\alpha}$,

$$\frac{dR_h}{dt} = R_h(N_Q + N_{t^c} + N_{H_2} + \tilde{\alpha}b). \quad (15)$$

where

$$\begin{aligned} N_Q &= 2C_Q\tilde{\alpha} - Y_h \\ N_{t^c} &= 2C_{t^c}\tilde{\alpha} - 2Y_h \\ N_{H_2} &= -3Y_h \end{aligned} \tag{16}$$

and the QCD Casimirs are $C_Q = C_{t^c} = 4/3$. The RGE in this case is thus:

$$\frac{dR_h}{dt} = \tilde{\alpha}R_h[(r_h + b) - 6R_h]. \tag{17}$$

where

$$r_h \equiv \sum_R 2C_R = 16/3. \tag{18}$$

and in the MSSM the QCD beta function is $b = -3$, which leads to the Pendleton-Ross fixed point $R_h^* = 7/18$.

In general we can always relabel the Yukawa couplings of any theory as h_i where $i = 1, \dots, n$ now runs over the n non-zero physically distinct couplings in the theory, and should not be confused with the previous use of the index i . Defining $Y_i \equiv h_i^2/(16\pi^2)$ and $R_i \equiv Y_i/\tilde{\alpha}$, we can always write the RG equations for the physically distinct Yukawa couplings of the theory as

$$\frac{dR_i}{dt} = \tilde{\alpha}R_i \left[(r_i + b) - \sum_j S_{ij}R_j \right], \tag{19}$$

where $r_i = \sum_R 2C_R$ with the sum being over the representations of the three fields associated with the trilinear coupling h_i , and S_{ij} is some matrix whose numerical value is fully specified by the wavefunction anomalous dimensions. The fixed point is then reached when the right hand side of Eq.19 is zero for all i . Thus, writing the fixed point solutions as R_i^* there are two fixed point values for each coupling: $R_i^* = 0$ or the solution corresponding to

$$[(r_i + b) - \sum_j S_{ij}R_j^*] = 0 \tag{20}$$

The non-trivial fixed point solution is:

$$R_i^* = \sum_{j=1}^n (S^{-1})_{ij}(r_j + b) \tag{21}$$

We shall begin our discussion by considering only the non-trivial fixed points, and later extend our analysis to the general case where some of the couplings are zero at the fixed point.

To determine the infra-red stability of the system in Eq.19, we need to Taylor expand it around the fixed point given in Eq.21. We can then drop all except the linear terms, the resulting system of which allows an algebraic solution and can thus

be tested for infra-red stability. We therefore make a change of variables to $\rho_i(t) \equiv R_i(t) - R_i^*$. The RGE Eq.19 then becomes

$$\frac{d\rho_i(t)}{dt} = \tilde{\alpha}(t)(\rho_i(t) + R_i^*) \left[(r_i + b_i) - \sum_{j=1}^n S_{ij}(\rho_j(t) + R_j^*) \right] \quad (22)$$

where we have substituted the fixed point values of R_i^* from Eq.21.

When we drop the quadratic terms in Eq.22 and change the independent variable from t to $\tilde{\alpha}$ by Eq.3, we obtain the linearised system

$$\frac{d\rho_i(t)}{d \ln \tilde{\alpha}(t)} \approx \frac{1}{b} R_i^* \sum_{j=1}^n S_{ij} \rho_j(t) \quad (23)$$

(no sum on i .) Eq.23 then describes the behaviour of the trajectories as they approach the fixed point. It has a general solution:

$$\rho_i(t) \approx \sum_{k=1}^n a_k x_i^{(k)} (\tilde{\alpha}(t))^{\lambda_k}, \quad (24)$$

where $x_j^{(k)}, \lambda_k$ are the $k = 1, \dots, n$ eigenvectors and eigenvalues of the eigenvalue equation

$$\begin{aligned} \sum_{j=1}^n A_{ij} x_j^{(k)} &= \lambda_k x_i^{(k)} \\ A_{ij} &\equiv \frac{1}{b} R_i^* S_{ij}. \end{aligned} \quad (25)$$

The a_k in the solutions are constants which describe the particular linear combination of eigenvectors and are set by initial boundary conditions. The infra-red stability properties of the solutions in Eq.24 are independent of the sign of the one-loop gauge beta function b , as we shall now show.

For $b > 0$, $\tilde{\alpha}$ decreases with decreasing renormalisation scale μ . We require every eigenvalue λ_k of the matrix $\frac{1}{|b|} R_i^* S_{ij}$ to have a positive real part for infra-red stability, so that $\rho_i \rightarrow 0$ as μ decreases. For $b < 0$, $\tilde{\alpha}$ increases with decreasing renormalisation scale μ , in this case we require all the eigenvalues of the matrix $-\frac{1}{|b|} R_i^* S_{ij}$ to have negative real parts, which is equivalent to the previous condition for $b > 0$, so the stability condition is independent of the sign of the beta function, as claimed. $\lambda_k = 0$ corresponds to a direction in coupling space which is neither attracted to, nor repelled by the fixed point (to first order). For each of these directions there should be one free parameter in the solution to the fixed point equations which embodies the information of where a solution lies along this direction (and is set by the initial boundary conditions). This corresponds to some information about the higher energy physics being retained at lower energies.

Complex eigenvalues always come in complex conjugate pairs, as do their associated eigenvectors. Writing $\lambda_k \equiv t_k + i s_k$, where t_k, s_k are real, the solution in this

case is

$$\rho_i \approx \sum_k a_k (x_i^{(k)} \tilde{\alpha}^{t_k + i s_k} + x_i^{(k)*} \tilde{\alpha}^{t_k - i s_k}) = \sum_k a_k \tilde{\alpha}^{t_j} [x_i^{(k)} \tilde{\alpha}^{i s_k} + x_i^{(k)*} \tilde{\alpha}^{-i s_k}] \quad (26)$$

where the sum runs over all values of k corresponding to pairs of complex eigenvalues. Eq.26 describes a spiral-like solution, the distance to the fixed point being controlled by $\tilde{\alpha}^{t_k}$. Thus t_k must be positive for the trajectory to be infra-red stable. For $b < 0$, $\tilde{\alpha}$ increases with decreasing normalisation scale so we require $Re(\lambda_k) < 0$ for all k for the solution to be stable in the infra-red direction. If any of the conditions are not met, the fixed point is either a saddle point or an ultra-violet fixed point and so the fixed point will never be achieved at low energies and must be rejected. In such cases we are led to consider other fixed points where some of the couplings take zero values at the fixed point. Another motivation for studying such fixed points would be if the non-trivial fixed point solution led to values of R_i with unphysical negative values (see example later).

As already remarked, in general for each value of i there are two possible fixed point solutions to consider: $R_i^* = 0$ or its non-zero fixed point value considered above. There are thus 2^n fixed points contained in the theory (some of which could be degenerate), and so far we have only considered one possibility (the non-zero case for each coupling). For each of the remaining possibilities it is straightforward to determine the fixed points. Suppose in general that we are considering a possibility with m zero fixed point solutions and $n - m$ non-zero solutions. First, we re-order the n couplings such that

$$\begin{aligned} [R_i^*]_{i=1,\dots,m} &= 0 \\ [R_i^*]_{i=m+1,\dots,n} &= \sum_{j=m+1}^n (S^{-1})_{ij} (r_j + b) \end{aligned} \quad (27)$$

Note that the non-zero couplings are now determined by the lower right hand $(n - m) \times (n - m)$ block of the re-ordered matrix S . As we shall see the discussion of the non-zero solutions turns out to be similar to that previously considered, while that of the zero fixed points is even simpler.

For the general case including m zero couplings, we need to Taylor expand around the fixed point given in Eq.27. We make a change of variables to $\rho_i(t) \equiv R_i(t) - R_i^*$, where $R_i^* = 0$ for $i = 1, \dots, m$. The RGE Eq.19 then becomes

$$\begin{aligned} \left[\frac{d\rho_i(t)}{dt} \right]_{i=1,\dots,m} &= \tilde{\alpha}(t) \rho_i(t) \left[(r_i + b) - \sum_{j=1}^m S_{ij} \rho_j(t) - \sum_{j=m+1}^n S_{ij} (\rho_j(t) + R_j^*) \right] \\ \left[\frac{d\rho_i(t)}{dt} \right]_{i=m+1,\dots,n} &= \tilde{\alpha}(t) (\rho_i(t) + R_i^*) \left[(r_i + b) - \sum_{j=1}^m S_{ij} \rho_j(t) - \sum_{j=m+1}^n S_{ij} (\rho_j(t) + R_j^*) \right] \end{aligned}$$

where we have substituted the fixed point values of R_i^* from Eq.27. When we drop the quadratic terms and change the independent variable from t to $\tilde{\alpha}$ by Eq.3, we

obtain the linearised system

$$\begin{aligned} \left[\frac{d\rho_i(t)}{d \ln \tilde{\alpha}(t)} \right]_{i=1,\dots,m} &\approx -\frac{\rho_i(t)}{b} \left[(r_i + b) - \sum_{j=m+1}^n S_{ij} R_j^* \right] \\ \left[\frac{d\rho_i(t)}{d \ln \tilde{\alpha}(t)} \right]_{i=m+1,\dots,n} &\approx \frac{1}{b} R_i^* \sum_{j=1}^n S_{ij} \rho_j(t) \end{aligned} \quad (28)$$

Eq.28 then describes the behaviour of the trajectories as they approach the fixed point.

For the $i = 1, \dots, m$ the equations are of the simple form:

$$\left[\frac{d\rho_i(t)}{d \ln \tilde{\alpha}(t)} \right]_{i=1,\dots,m} \approx \lambda_i \rho_i(t) \quad (29)$$

where

$$\lambda_i = \frac{1}{b} \left[\sum_{j=m+1}^n S_{ij} R_j^* - (r_i + b) \right] \quad (30)$$

with solution,

$$[\rho_i(t)]_{i=1,\dots,m} = a_i(\tilde{\alpha})^{\lambda_i} \quad (31)$$

As in the previous argument, for $b > 0$, in order to have $\rho_i \rightarrow 0$ in the infra-red we require $\lambda_i > 0$, where here λ_i is given by Eq.30.

For $i = m + 1, \dots, n$, assuming that the m zero solutions are infra-red stable, as discussed in the above paragraph, we may simplify the procedure by working in the infra-red region where $[\rho_i]_{i=1,\dots,m} \rightarrow 0$, so that we have a simplified set of equations to solve:

$$\left[\frac{d\rho_i(t)}{d \ln \tilde{\alpha}(t)} \right]_{i=m+1,\dots,n} \approx \frac{1}{b} R_i^* \sum_{j=m+1}^n S_{ij} \rho_j(t) \quad (32)$$

In fact the argument now parallels that given previously for the non-zero solutions, as in Eq.23 and the discussion below it, with attention now being focussed on the $(n - m) \times (n - m)$ lower right hand block of the re-ordered matrix S .

As a simple example of the application of these results, we consider the example of the next-to-minimal supersymmetric standard model (NMSSM) described by the superpotential:

$$W = h Q t^c H_2 + \lambda N H_1 H_2 - \frac{k}{3} N^3 \quad (33)$$

where we have included in addition to the top quark Yukawa coupling the NMSSM terms which contain the gauge singlet N . The fixed points of this theory in the presence of the gauge couplings have been previously studied numerically [11]. It was found there that the couplings λ and k are attracted towards zero values [11], and it is interesting to see how our analytic treatment here reproduces this. The relevant

RGEs are [12, 11], keeping only the QCD gauge coupling:

$$\begin{aligned}
16\pi^2 \frac{dk}{d\ln\mu} &= k(6k^2 + 6\lambda^2) \\
16\pi^2 \frac{d\lambda}{d\ln\mu} &= \lambda(2k^2 + 4\lambda^2 + 3h^2) \\
16\pi^2 \frac{dh}{d\ln\mu} &= h(\lambda^2 + 6h^2 - 2g^2(C_Q + C_{t^c}))
\end{aligned} \tag{34}$$

In the notation of Eqs.9, 11 we find:

$$\begin{aligned}
\frac{dR_k}{dt} &= \tilde{\alpha}R_k[b - 6R_k - 6R_\lambda] \\
\frac{dR_\lambda}{dt} &= \tilde{\alpha}R_\lambda[b - 2R_k - 4R_\lambda - 3R_h] \\
\frac{dR_h}{dt} &= \tilde{\alpha}R_h[(r_h + b) - R_\lambda - 6R_h]
\end{aligned} \tag{35}$$

where $r_h = 2(C_Q + C_{t^c})$.

We now order the ratios as $R_i = (R_k, R_\lambda, R_h)$ and write the RGEs for the ratios in the form of Eq.19 where we identify:

$$r_1 = 0, \quad r_2 = 0, \quad r_3 = r_h = 16/3 \tag{36}$$

with $b = -3$ for the NMSSM as in the MSSM, and the matrix S given by:

$$S = \begin{pmatrix} 6 & 6 & 0 \\ 2 & 4 & 3 \\ 0 & 1 & 6 \end{pmatrix} \tag{37}$$

The non-zero couplings are obtained by inverting the matrix S , then using Eq.21 with $(r_j + b) = (-3, -3, 7/3)$. The result is:

$$R_i^* = \left(\frac{29}{18}, -\frac{19}{9}, \frac{20}{27} \right) \tag{38}$$

The negative value of the fixed point for R_λ means that we must abandon this fixed point as a physical possibility.

We next try a fixed point with $R_\lambda^* = 0$, and the other two couplings being given by their non-zero solutions. Following our procedure we first re-order the couplings so that those with zero fixed point values are listed first. In this case we re-order the couplings as : $R_i = (R_\lambda, R_k, R_h)$, with $(r_j + b) = (-3, -3, 7/3)$ (unchanged), but now we have:

$$S = \begin{pmatrix} 4 & 2 & 3 \\ 6 & 6 & 0 \\ 1 & 0 & 6 \end{pmatrix} \tag{39}$$

Since by fiat $R_1^* = 0$ it only remains to determine the fixed point values for the other two couplings. Since the lower 2×2 block of the re-ordered matrix S , is proportional to the unit matrix in this case it is easy to invert and we find

$$(R_k^*, R_h^*) = \left(-\frac{1}{2}, \frac{7}{18}\right) \quad (40)$$

For $R_k^* = 0$ with the others non-zero, we find

$$(R_\lambda^*, R_h^*) = \left(-\frac{25}{21}, \frac{37}{63}\right). \quad (41)$$

The trial fixed point in these cases again contains a non-physical negative value, so we abandon these possibilities.

Finally we try a fixed point with $R_\lambda = R_k = 0$, but with $R_h \neq 0$. No further re-ordering is required since the two zero ratios occur for $i = 1, 2$. The non-zero fixed point for the top Yukawa coupling is given by:

$$R_h^* = R_3^* = S_{33}^{-1}(r_3 + b) = \frac{7}{18} \quad (42)$$

Having obtained a physically sensible fixed point we must now test its infra-red stability. The behaviour of the couplings $[R_i]_{i=1,2}$ around the origin is determined by the eigenvalues in Eq.30 which are given by:

$$[\lambda_i]_{i=1,2} = \frac{1}{b}[S_{i3}R_3^* - (r_i + b)] = [-1, -\frac{25}{18}] \quad (43)$$

which indicates that this fixed point is attractive in the infra-red direction. The behaviour of the coupling R_3 around its fixed point is governed by the single entry matrix

$$A_{33} = \frac{1}{b}R_3^*S_{33} = -\frac{7}{9} \quad (44)$$

whose eigenvalue is trivially equal to $\lambda = -\frac{7}{9}$. Since $b < 0$ in this case, a negative eigenvalue indicates an IRSFP. In fact $R_h^* = 7/18$ is just the usual Pendleton-Ross fixed point of the MSSM for the top quark Yukawa coupling discussed previously.

There are several important technical questions not addressed by the above arguments. One of these questions is: how fast [13] does the solution approach the fixed point? So far, we can only suggest a numerical test where the boundary conditions are picked from some range of initial parameters spanning a range centred upon 1 and of the order 1. We know that the IRSFP is only formally realised in the limit $\mu \rightarrow 0$. While it is clear that when the magnitude of λ_i is large, so is the speed of the running and therefore the rate at which the fixed point is approached, in models of particle physics μ never reaches this limit and so the fixed point solution can only be used approximately. Some measure on the “error” of this prediction would then be useful. It is not clear yet if there are general classes of models which possess an IRSFP. Ideally, there would only be one IRSFP in the theory. If two exist in the same model, it would be unclear a priori which would be more likely to be realised in

nature. Finally, we should note that the solutions $R_{i>m}^*$ to Eq.27 are not all guaranteed to be positive. Any that are negative will never be reached: the coupling would simply approach zero and therefore should be added to the list of couplings $R_{i=1,\dots,m}$.

To summarise, we have determined the fixed points in a large class of supersymmetric models. These fixed points offer the possibility of increased predictivity and therefore testability of the models. There are however there are several important theoretical questions raised by this approach. Perhaps the most important is that of the existence and uniqueness of IRSFPs in any given theory. In the absence of such general theorems we have provided a well defined procedure for exploring the fixed point properties. Our suggested procedure is the following: start with the case of a non-trivial fixed point for all the n physical couplings, and examine its stability; then go on to the case with one zero fixed point coupling and $n - 1$ non-zero fixed point couplings and examine its stability; and so on down to the case of the completely trivial fixed point with n zero fixed point couplings. We have the given conditions for infra-red stability for the general case of m zero fixed point couplings and $n - m$ non-zero fixed point couplings, which enables this procedure to be followed. Applying the analysis to the NMSSM in the low $\tan\beta$ regime, we are able to show that the fixed point prediction for the top quark mass is equivalent to the MSSM prediction. These techniques may also be applied to the fermion mass problem [10].

References

- [1] B. Pendleton and G.G. Ross, Phys. Lett. **B98** (1981) 21.
- [2] C.T. Hill, Phys. Rev. **D24** (1981) 691.
- [3] M. Lanzagorta and G.G. Ross, Phys. Lett. **B349** (1995) 319
- [4] G.G. Ross, Phys. Lett. **B364** (1995) 216.
- [5] B. Schrempp, Phys. Lett. **B344** (1995) 193-200.
- [6] M. Lanzagorta and G.G. Ross, Phys. Lett. **B364** (1995) 163.
- [7] I. Jack and D.R.T. Jones, Phys. Lett. **B349** (1995) 294;
P.M. Ferreira, I. Jack and D.R.T. Jones, Phys. Lett. **B357** (1995) 359;
P.M. Ferreira, I. Jack and D.R.T. Jones, Phys. Lett. **B392** (1997) 276.
- [8] B.C. Allanach, G. Amelino-Camelia and O. Phillipsen, Phys. Lett. **B393**
- [9] S.P. Martin and M.T. Vaughn, Phys. Rev. **D 50** (1994) 2282.
- [10] B.C. Allanach and S.F. King, RAL 97-007, SHEP 97-01, hep-ph/9703293. (1997) 349.
- [11] T. Elliott, S.F.King and P.L.White, Phys. Rev. **D49** (1994) 2435.
- [12] B.C.Allanach and S.F.King, Phys. Lett. **B328** (1994) 360.
- [13] M. Bando, J. Sato and K. Yoshioka, ICRR Report-384-97-7, KUNS 1437 HE(TH)97/04, hep-ph/9703321.